

CONSTANTS:

Avogadro's number: $N_A = 6.02 \times 10^{23}$

Boltzmann's constant: $k = 1.38 \times 10^{-23}$ J/K

Universal gas constant: $R = 0.0821$ L·atm/mol·K

Universal gas constant: $R = 8.314$ J/mol·K

Atomic mass unit: $u = 1.66 \times 10^{-27}$ kg

EQUATIONS:

$$PV = nRT$$

$$PV = NkT$$

$$\frac{3}{2}kT = \frac{1}{2}m\overline{v^2} \quad T_C = T_K + 273^\circ$$

1. According to the kinetic theory of gases, the temperature of an ideal gas is directly proportional to the
 - (a) volume of the gas.
 - (b) mean distance between collisions between particles.
 - (c) angular momentum of the particles.
 - (d) average kinetic energy of the particles.
 - (e) average momentum of the particles.

2. The hydrogen molecules in a container have the same root-mean-square speed as the oxygen molecules in another container. Which of the following conclusions can be made with certainty?
 - (a) the oxygen gas will have the higher temperature.
 - (b) the hydrogen gas will have the higher temperature.
 - (c) both gases have the same temperature.
 - (d) the hydrogen gas has the higher pressure.
 - (e) both gases have the same pressure.

3. Which of the following is a notable failure of the ideal gas model?
 - (a) the condensation of gases
 - (b) the expansion of gases as they warm
 - (c) the relationship between temperature and molecular kinetic energy
 - (d) the proportionality of pressure and temperature

4. If you keep the volume of a sample of gas constant while its temperature is allowed to change,
 - (a) the pressure of the gas will remain constant while the temperature increases.
 - (b) the root-mean-square speed of its particles will remain constant.
 - (c) the pressure will decrease as the temperature increases.
 - (d) the pressure will be directly proportional to the Kelvin temperature.

5. If the temperature of an ideal gas is kept constant, while its pressure and volume are permitted to change
 - (a) its volume will vary directly as the pressure.
 - (b) the product of its pressure and volume will remain constant.
 - (c) its pressure will remain constant while its volume varies.
 - (d) its volume will remain constant while its pressure varies.


6. A sample of oxygen gas and a sample of hydrogen gas are stored in the same store room at the same temperature. The mass of a molecule of oxygen is 32 u. The mass of a hydrogen molecule is 2 u. The ratio of the average kinetic energy of the oxygen molecules to that of the hydrogen molecules is
 - (a) 1 to 1.
 - (b) 4 to 1.
 - (c) 16 to 1.
 - (d) 1 to 16.

7. A sample of oxygen gas and a sample of hydrogen gas are stored in the same store room at the same temperature. The mass of a molecule of oxygen is 32 u. The mass of a hydrogen molecule is 2 u. The ratio of v_{rms} of the oxygen molecules to that of the hydrogen molecules is
 - (a) 1 to 1.
 - (b) 4 to 1.
 - (c) 1 to 4.
 - (d) 1 to 16.


8. The temperature of a gas is 10°C . To double the average kinetic energy of its molecules, the temperature of the gas must be raised to
 - (a) 20°C .
 - (b) 40°C .
 - (c) 293°C .
 - (d) 566°C .

9. Which of the green plots best represents the distribution of molecular speeds in a gas at 500 K if the purple curve represents this distribution for the same gas at 300 K?


(a)




(b)




(c)



(d)



(e)



1. Find v_{rms} for Ar gas at 20°C . [428 m/s]
2. Find the ratio of v_{rms} for O_2 and H_2 at the same temperature. [1:4]
3. (a) What is the average kinetic energy for nitrogen molecules, N_2 , at 20°C ? [Note: nitrogen *atoms* are denoted as ${}^{14}_7\text{N}$.]

$$\underline{6.1 \times 10^{-21} \text{ J}}$$

- (b) What is the root-mean-square speed for these nitrogen molecules?

$$\underline{500 \text{ m/s}}$$

4. A 0.02 m^3 sample of a gas at a pressure of 1000. kPa is allowed to expand at constant temperature until its pressure decreases to 500 kPa. What will the new volume of the gas be?

$$\underline{0.04 \text{ m}^3}$$